

Supplementary

Supplementary table 1 Demographics for patients who had their surgery postponed due to the COVID-19 pandemic.

	Knee arthroplasty (included) n = 59	Knee arthroplasty (drop-out) n = 43	p-value	Hip arthroplasty (included) n = 51	Hip arthroplasty (drop-out) n = 23	p-value
Age, yrs, median (2.5-97.5 quantiles)	70 (62; 75)	69 (62; 75)	0.79	72 (61; 76)	70 (60; 75)	0.65
Sex, female, % (n)	58% (34)	51% (22)	0.48	67% (34)	74% (17)	0.49
BMI, kg/m ² , median (2.5-97.5 quantiles)	30 (26; 35)	29 (25; 33)	0.48	27 (24; 31)	27 (24; 34)	0.59
OXS, median (2.5-97.5 quantiles)	19 (13; 23)	19 (14; 24)	0.81	-	-	-
OHS, median (2.5-97.5 quantiles)	-	-	-	18 (12; 22)	16 (10; 23)	0.89
FJS, median (2.5-97.5 quantiles)	8 (2; 14)	8 (3; 25)	0.16	7 (0; 14)	3 (2; 13)	0.72
EQ5D index, median (2.5-97.5 quantiles)	0.66 (0.34; 0.72)	0.66 (0.34; 0.72)	0.49	0.59 (0.32; 0.66)	0.39 (0.24; 0.66)	0.24
EQ5D VAS, median (2.5-97.5 quantiles)	55 (38; 70)	50 (30; 74)	0.94	45 (21; 70)	55 (28; 70)	0.63
Pain VAS, median (2.5-97.5 quantiles)	72 (62; 82)	75 (66; 85)	0.64	78 (71; 82)	75 (64; 87)	0.67
ASA classification, % (n) ^b						
1	5% (3)	2% (1)		10% (5)	9% (2)	
2	80% (47)	51% (22)		61% (31)	52% (12)	
3	15% (9)	5% (2)	-	29% (15)	30% (7)	-
4	-	2% (1)		-	-	
Kellgren & Lawrence Classification ^a , % (n)						
2	2% (1)	6% (3)		-	-	
3	42% (25)	47% (20)	0.36	-	-	-
4	56% (33)	47% (20)		-	-	
Tönnis Classification ^a , % (n) ^c						
2	-	-		20% (10)	22% (5)	
3	-	-	-	78% (40)	78% (18)	0.99

ASA = American Society of Anesthesiologists; EQ5D = EuroQol 5-Dimension; FJS = Forgotten Joint Score; OHS = Oxford Hip Score; OXS = Oxford Knee Score; VAS = visual analogue scale.

a) For bilateral surgery, the side with the highest score was used.

b) 17 and 2 missing in knee and hip replacement (drop-out) groups, respectively

c) 1 missing in hip replacement (included) group.